

# Bolivia 2012 (Ages 13-15) Global Youth Tobacco Survey (GYTS)



## FACT SHEET . . . . .

The Bolivia GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Bolivia could include in a comprehensive tobacco control program.

The Bolivia GYTS was a school-based survey of students in grade 6<sup>th</sup> of primary and 1<sup>st</sup> – 5<sup>th</sup> in secondary conducted in 2012.

A two-stage cluster sample design was used to produce representative data for Bolivia. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. A total of 3,373 students participated in the Bolivia GYTS of which 2,263 were age 13 to 15 years. The overall response rate of all students surveyed was 88.9%.

### Prevalence

- 31.9% of students had ever smoked cigarettes (Boys = 40.1%, Girls = 28.2%)
- 18.7% currently use any tobacco product (Boys = 20.9%, Girls = 16.4%)
- 11.3% currently smoke cigarettes (Boys = 15.3%, Girls = 9.9%)
- 10.4% currently use other tobacco products (Boys = 9.9%, Girls = 9.6%)
- 20.6% of never smokers are likely to initiate smoking next year (Boys = 36.5%, Girls = 14.7%)

### Knowledge and Attitudes

- 25.1% think boys and 24.7% think girls who smoke have more friends
- 16.9% think boys and 12.6% think girls who smoke look more attractive

### Access and Availability - Current Smokers

- 13.1% usually smoke at home
- 51.1% buy cigarettes in a store
- 90.4% who bought cigarettes in a store were NOT refused purchase because of their age

### Exposure to Secondhand Smoke (SHS)

- 34.5% live in homes where others smoke in their presence
- 46.6% are around others who smoke in enclosed public places
- 50.3% are around others who smoke in outdoor public places
- 86.5% think smoking should be banned from public places
- 63.5% think smoke from others is harmful to them
- 41.3% have one or more parents who smoke
- 9.7% have most or all friends who smoke

### Cessation - Current Smokers

- 61.6% want to stop smoking
- 68.6% tried to stop smoking during the past year
- 65.7% have ever received help to stop smoking

### Media and Advertising

- 71.9% saw anti-smoking media messages, in the past 30 days
- 65.4% saw pro-cigarette ads on billboards, in the past 30 days
- 53.0% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 11.7% have an object with a cigarette brand logo
- 13.6% were offered free cigarettes by a tobacco company representative

### School

- 49.1% had been taught in class, during the past year, about the dangers of smoking
- 24.1% had discussed in class, during the past year, reasons why people their age smoke
- 51.0% had been taught in class, during the past year, the effects of tobacco use

### Highlights

- 18.7% of students currently use any form of tobacco; 11.3% currently smoke cigarettes; 10.4% currently use some other form of tobacco
- SHS exposure – more than one-third of the students live in homes where others smoke, and 46.6% of students are around others who smoke in enclosed public places; two in five students have at least one parent who smokes
- More than three in five students think smoke from others is harmful to them
- 86.5% of students think smoking in public places should be banned
- Over six in 10 current smokers want to stop smoking
- 11.7% of students have an object with a cigarette brand logo on it
- 13.6% were offered free cigarettes by a tobacco company representative
- More than seven in 10 students saw anti-smoking media messages in the past 30 days; nearly two-thirds of the students saw pro-cigarette ads on billboards and over one-half saw pro-cigarette ads in newspapers or magazines in the past 30 days