

Montenegro (Ages 13-15)

Global Youth Tobacco Survey (GYTS)

FACT SHEET



The Montenegro GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Montenegro could include in a comprehensive tobacco control program.

The Montenegro GYTS was a school-based survey of students in grades VI, VII, VIII and I conducted in 2008.

A two-stage cluster sample design was used to produce representative data for Montenegro. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the student response rate was 93.2%, the class response rate was 99.7% and the overall response rate was 92.9%. A total of 5,723 students participated in the Montenegro GYTS of which 3,299 were aged 13 to 15 years.

Prevalence

- 31.3% of students had ever smoked cigarettes (Boys = 30.7%, Girls = 31.9%)
 - 6.3% currently use any tobacco product (Boys = 6.6%, Girls = 5.9%)
 - 5.1% currently smoke cigarettes (Boys = 5.7%, Girls = 4.4%)
 - 1.0% currently daily cigarette smokers (Boys = 0.9%, Girls = 0.9%)
 - 3.6% used any form of tobacco other than cigarettes in the past 30 days (Boys = 3.7%, Girls = 3.5%)
- 39.6% ever smokers initiated smoking before age ten (Boys = 40.6%, Girls = 38.8%)
- 16.0% of never smokers are likely to initiate smoking next year (Boys = 15.7%, Girls = 16.5%)

Access and Availability - Current Smokers

- 23.7% current smokers who usually smoke at home
- 55.6% buy cigarettes in a store
- 96.5% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

- 76.8% live in homes where others smoke in their presence
- 69.9% are around others who smoke in places outside their home
- 89.8% think smoking should be banned from public places
- 67.4% think smoke from others is harmful to them
- 60.3% have one or more parents who smoke
- 11.5% have most or all friends who smoke

Cessation - Current Smokers

- 41.2% want to stop smoking
- 75.6% tried to stop smoking during the past year
- 65.0% have ever received help to stop smoking
 - 7.9% always have or feel like having a cigarette first thing in the morning

Media and Advertising

- 89.8% saw anti-smoking media messages in the past 30 days
- 47.4% saw pro-cigarette ads on billboards in the past 30 days
- 58.2% saw pro-cigarette ads in newspapers or magazines in the past 30 days
- 20.2% have an object with a cigarette brand logo
- 19.3% were offered free cigarettes by a tobacco company representative

School

- 56.0% had been taught in class, during the past year, about the dangers of smoking
- 41.2% had discussed in class, during the past year, reasons why people their age smoke
- 52.5% had been taught in class, during the past year, the effects of tobacco use

Highlights

- Less than 1 in 10 students currently use any form of tobacco; 5.1% currently smoke cigarettes.
- SHS exposure is HIGH – over 7 in 10 students live in homes where others smoke, nearly 7 in 10 students are around others who smoke in places outside of their home; 6 in 10 students have one or more parent who smoke.
- Nearly 9 in 10 students think smoking should be banned from public places.
- Over 4 in 10 students who are currently smoking indicated that they want to stop smoking now; more than 7 in 10 students currently smoking tried to stop during the past year.
- Nearly 9 in 10 students saw anti-smoking media messages while about 50% saw pro-cigarette ads on billboards, and approximately 6 in 10 students saw pro-cigarette ads in newspapers & magazines.