GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:

mpower

Monitor tobacco use & prevention policies

Protect people from tobacco smoke

Offer help to quit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising, promotion, & sponsorship

Raise taxes on tobacco

GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco advertising and promotion, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is selfadministered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In Poland, GYTS was conducted in 2022 by the National Research Institute of Oncology, under the coordination of the Ministry of Health. The overall response rate was 74.8%. A total of 3,985 eligible students in 7th and 8th grade in elementary school, and 1st grade in high school completed the survey, of which 3,573 were aged 13-15 years. Data are reported for students aged 13-15 years.

GYTS Highlights

TOBACCO USE

- 17.1% of students, 17.3% of boys, and 16.9% of girls currently used any tobacco products (includes heated tobacco products).
- 12.5% of students, 12.2% of boys, and 12.7% of girls currently smoked tobacco.
- 11.7% of students, 11.2% of boys, and 12.3% of girls currently smoked cigarettes.
- 10.0% of students, 9.7% of boys, and 10.4% of girls currently used heated tobacco products.
- 4.5% of students, 5.0% of boys, and 4.0% of girls currently used smokeless tobacco.

ELECTRONIC CIGARETTES

• 22.3% of students, 21.2% of boys, and 23.4% of girls currently used electronic cigarettes.

CESSATION

- More than 5 in 10 (54.4%) students who currently smoked tobacco tried to stop smoking in the past 12 months.
- Almost 3 in 10 (27.8%) students who currently smoked tobacco wanted to stop smoking now.

SECONDHAND SMOKE

- 28.8% of students were exposed to tobacco smoke at home.
- 41.9% of students were exposed to tobacco smoke inside enclosed public places.

ACCESS & AVAILABILITY

- 44.1% of students who currently smoked cigarettes bought cigarettes from a shop, supermarket, street vendor, or kiosk.
- Among students who currently smoked cigarettes who tried to buy cigarettes, 64.1% were not prevented from buying them because of their age.

ADVERTISING & PROMOTION

- More than 3 in 10 (34.9%) students noticed anti-tobacco messages in the media.
- Almost 3 in 10 (29.9%) students noticed tobacco advertisements or promotions when visiting points of sale.
- Almost 1 in 10 (8.6%) students had something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 34.4% of students definitely thought other people's tobacco smoking is harmful to them.
- 69.6% of students favored prohibiting smoking inside enclosed public places.





Ministry of Health Republic of Poland






The mark "CDC" is owned by the US Dept. of Health and Human Services and is used with permission. Use of this logo is not an endorsement by HHS or CDC of any particular product, service, or enterprise.


GYTS GLOBAL YOUTH TOBACCO SURVEY

FACT SHEET POLAND 2022

TOBACCO USE

10011000 002			
ANY TOBACCO USE (smoked, heated, and/or smokeless)	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco users ¹	17.1	17.3	16.9
Ever tobacco users ²	41.6	42.2	41.0
SMOKED TOBACCO			
Current tobacco smokers ³	12.5	12.2	12.7
Current cigarette smokers ⁴	11.7	11.2	12.3
Frequent cigarette smokers ⁵	3.5	3.2	4.0
Current smokers of other tobacco ⁶	5.3	5.9	4.7
Ever tobacco smokers ⁷	35.0	34.8	35.3
Ever cigarette smokers ⁸	30.7	29.9	31.6
Ever smokers of other tobacco ⁹	16.1	18.5	13.6*
HEATED TOBACCO			
Current users of heated tobacco products ¹⁰	10.0	9.7	10.4
Ever users of heated tobacco products ¹¹	20.9	20.6	21.3
SMOKELESS TOBACCO			
Current smokeless tobacco users ¹²	4.5	5.0	4.0
Ever smokeless tobacco users13	13.3	14.6	11.9
SUSCEPTIBILITY			
Never to bacco users susceptible to to bacco use in the future $^{\rm 14}$	39.3	34.5	44.3*
Never tobacco smokers who thought they might enjoy smoking a cigarette ¹⁵	20.2	19.6	20.7





ELECTRONIC CIGARETTES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current electronic cigarette users ¹⁶	22.3	21.2	23.4
Ever electronic cigarette users ¹⁷	41.2	42.2	40.1
CESSATION			
	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers who tried to stop smoking in the past 12 months	54.4	51.7	57.0
Current tobacco smokers who wanted to stop smoking now	27.8	28.1	27.2
Current tobacco smokers who thought they would be able to stop smoking if they wanted to	79.0	83.9	74.3*
Current tobacco smokers who have ever received help/advice from a program or professional to stop	7.6	7.3	7.9

SECONDHAND SMOKE

smoking

	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home ¹⁸	28.8	26.4	31.5*
Exposure to tobacco smoke inside any enclosed public place ¹⁸	41.9	37.2	47.0*
Exposure to tobacco smoke at any outdoor public $place^{18}$	59.0	51.8	66.9*
Students who saw anyone smoking inside the school building or outside on school property ¹⁹	51.3	49.3	53.6

ACCESS & AVAILABILITY

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current cigarette smokers who bought cigarettes from a shop, supermarket, street vendor, or kiosk ²⁰	44.1	48.7	40.5
Current cigarette smokers who were not prevented from buying cigarettes because of their age 21	64.1	53.9	74.4*
Current cigarette smokers who bought cigarettes as individual sticks ²²	19.4	20.9	18.3
ADVERTISING & PROMOTION			
TOBACCO ADVERTISING & PROMOTION	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who noticed tobacco advertisements or promotions at points of sale ²³	29.9	30.3	29.2
Students who saw anyone using tobacco on television, videos, or movies ²⁴	61.9	62.2	61.5
Students who were ever offered a free tobacco product from a tobacco company representative	6.8	8.2	5.2*
Students who had something with a tobacco brand logo on it	8.6	9.3	7.7
ANTI-TOBACCO ADVERTISING & PROMOTION			
Students who noticed anti-tobacco messages in the media ¹⁹	34.9	34.5	35.1
Students who noticed anti-tobacco messages at sporting or community events ²⁵	30.9	31.7	29.7
Current tobacco smokers who thought about quitting because of a warning label ²⁶	15.6	17.4	14.0
Students who were taught in school about the dangers of tobacco use in the past 12 months	43.1	42.3	44.1
KNOWI EDGE & ATTITUDES			

KNOWLEDGE & ATTITUDES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	31.5	30.1	33.1
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	45.5	46.2	44.8
Students who definitely thought other people's tobacco smoking is harmful to them	34.4	37.6	30.7*
Students who favored prohibiting smoking inside enclosed public places	69.6	67.1	72.1*
Students who favored prohibiting smoking at outdoor public places	45.3	47.7	42.7*

¹ Smoked cigarettes, smoked other type of tobacco, and/or used smokeless tobacco anytime during the past 30 days. ² Ever smoked tobacco and/or used smokeless tobacco. ³ Smoked cigarettes or other type of tobacco anytime during the past 30 days. ⁴ Smoked cigarettes anytime during the past 30 days. ⁵ Smoked cigarettes anytime during the past 30 days. ⁵ Smoked cigarettes or other type of tobacco other than cigarettes anytime during the past 30 days. ⁷ Ever smoked cigarettes or other type of tobacco, even one or two puffs. ⁸ Ever smoked cigarettes or other type of tobacco, even one or two puffs. ¹⁰ Used heated tobacco orducts anytime during the past 30 days. ¹¹ Ever used heated tobacco products. ¹² Used smokeless tobacco anytime during the past 30 days. ¹³ Ever used smokeless tobacco. ¹⁴ Susceptible to future tobacco use includes those who answered "definitely yes", "probably yes", or "probably not" to using tobacco for of their best friends offerred it to them, or "definitely yes", "probably yes", or "probably not" to using tobacco ince is assessed separately from cigarettes, other smoked tobacco products, and heated tobacco products, and heated tobacco products, and heated tobacco use includes those who robacco during the past 30 days. ¹⁵ Ever used electronic cigarettes is assessed separately from cigarettes, other smoked tobacco products, and heated tobacco products, and heated tobacco products, and heated tobacco products and is not included in the current definition of current any tobacco use. ¹⁷ Ever used electronic cigarettes during the past 30 days. ²³ Among those who visited a point of sale in the past 30 days. ²⁴ Among those who watched television, videos, or movies in the past 30 days. ²⁵ Among those who attended sporting or community events in the past 30 days. ²⁶ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁶ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁶ Among those who noticed warn

NOTE: Data are weighted to be nationally representative of all students aged 13-15 years who are enrolled in school. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.

*Gender comparisons are statistically significant at p < 0.05.