GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:



Monitor tobacco use & prevention policies

Protect people from tobacco smoke Offer help to quit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising,

promotion, & sponsorship

Raise taxes on tobacco

GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly, and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco advertising and promotion, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is self-administered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In the Kingdom of Saudi Arabia, GYTS was conducted in 2022 by the Ministry of Health and the Ministry of Education. The overall response rate was 92.3%. A total of 6,983 eligible students in grades 1-3 intermediate completed the survey, of which 5,610 were aged 13-15 years. Data are reported for students aged 13-15 years.

GYTS Highlights

TOBACCO USE

- 9.4% of students, 10.6% of boys, and 8.0% of girls currently used any tobacco products (including shisha and heated tobacco products).
- 6.8% of students, 7.6% of boys, and 5.7% of girls currently smoked tobacco (including shisha).
- 2.9% of students, 3.5% of boys, and 2.2% of girls currently smoked cigarettes.
- 2.9% of students, 3.4% of boys, and 2.4% of girls currently smoked shisha.
- 1.5% of students, 1.6% of boys, and 1.4% of girls currently used heated tobacco products.
- 3.6% of students, 4.4% of boys, and 2.6% of girls currently used smokeless tobacco.

ELECTRONIC CIGARETTES

• 5.4% of students, 6.1% of boys, and 4.7% of girls currently used electronic cigarettes.

CESSATION

- More than 6 in 10 (64.9%) students who currently smoked tobacco tried to stop smoking in the past 12 months.
- Almost 6 in 10 (55.8%) students who currently smoked tobacco wanted to stop smoking now.

SECONDHAND SMOKE

- 18.1% of students were exposed to tobacco smoke at home.
- 27.2% of students were exposed to tobacco smoke inside enclosed public places.

ACCESS & AVAILABILITY

- 49.2% of students who currently smoked cigarettes bought cigarettes from a store, shop, street vendor, or the internet.
- Among students who currently smoked cigarettes who tried to buy cigarettes, 51.7% were not prevented from buying them because of their age.

ADVERTISING & PROMOTION

- More than 4 in 10 (43.4%) students noticed anti-tobacco messages in the media.
- Almost 2 in 10 (19.7%) students noticed tobacco advertisements or promotions when visiting points of sale.
- Almost 1 in 10 (9.3%) students had something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 50.3% of students definitely thought other people's tobacco smoking is harmful to them.
- 83.8% of students favored prohibiting smoking inside enclosed public places.







TOBACCO USE			
ANY TOBACCO USE (smoked, heated, and/or smokeless)	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco users ¹	9.4	10.6	8.0*
Ever tobacco users ²	31.1	34.3	27.6*
SMOKED TOBACCO			
Current tobacco smokers ³	6.8	7.6	5.7
Current cigarette smokers ⁴	2.9	3.5	2.2*
Frequent cigarette smokers ⁵	0.4	0.6	0.2*
Current smokers of other tobacco ⁶	5.7	6.2	5.0
Current smokers of shisha ⁷	2.9	3.4	2.4
Ever tobacco smokers ⁸	23.8	26.2	21.2*
Ever cigarette smokers ⁹	12.6	14.9	10.3*
Ever smokers of other tobacco ¹⁰	18.8	20.5	17.0
Ever smokers of shisha ¹¹	15.0	15.7	14.2
HEATED TOBACCO	4.5	4.6	
Current users of heated tobacco products ¹²	1.5 7.5	1.6 8.9	1.4 5.9*
Ever users of heated tobacco products ¹³	7.5	0.9	3.9
SMOKELESS TOBACCO	0.4		0.44
Current smokeless tobacco users ¹⁴	3.6	10.9	2.6*
Ever smokeless tobacco users ¹⁵	9.1	10.9	7.2*
SUSCEPTIBILITY			
Never to bacco users susceptible to to bacco use in the future 16	7.4	7.6	7.3
Never tobacco smokers who thought they might enjoy smoking a cigarette ¹⁷	18.3	21.2	15.6*
ELECTRONIC CIGARETTES			
	OVERALL (%)	BOYS (%)	GIRLS (%)
Current electronic cigarette users ¹⁸	5.4	6.1	4.7
Ever electronic cigarette users ¹⁹	14.4	15.8	13.0
CESSATION			
	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers who tried to stop smoking in the past 12 months	64.9	69.0	61.9
Current tobacco smokers who wanted to stop smoking now	55.8	59.7	52.2
Current tobacco smokers who thought they would be able to stop smoking if they wanted to	64.5	65.2	64.0
	64.5 21.4	65.2 22.1	64.0
would be able to stop smoking if they wanted to Current tobacco smokers who have ever received help/advice from a program or			
would be able to stop smoking if they wanted to Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking			
would be able to stop smoking if they wanted to Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking	21.4	22.1	18.4
would be able to stop smoking if they wanted to Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking SECONDHAND SMOKE	21.4 OVERALL (%)	22.1 BOYS (%)	18.4 GIRLS (%)
would be able to stop smoking if they wanted to Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking SECONDHAND SMOKE Exposure to tobacco smoke at home ²⁰ Exposure to tobacco smoke inside any enclosed	21.4 OVERALL (%) 18.1	22.1 BOYS (%) 14.4	18.4 GIRLS (%) 21.9*

ACCESS & AVAILABILITY					
	OVERALL (%)	BOYS (%)	GIRLS (%)		
Current cigarette smokers who bought cigarettes from a store, shop, street vendor, or the internet ²²	49.2	55.2	40.2		
Current cigarette smokers who were not prevented from buying cigarettes because of their age 23	51.7	52.7			
Current cigarette smokers who bought cigarettes as individual sticks ²⁴	30.9	29.1			
ADVERTISING & PROMOTION					
TOBACCO ADVERTISING & PROMOTION	OVERALL (%)	BOYS (%)	GIRLS (%)		
Students who noticed tobacco advertisements or promotions at points of sale ²⁵	19.7	19.6	20.0		
Students who saw anyone using tobacco on television, videos, or movies ²⁶	48.1	45.8	50.4		
Students who were ever offered a free tobacco product from a tobacco company representative	6.7	9.0	4.5*		
Students who had something with a tobacco brand logo on it	9.3	11.5	6.9*		
ANTI-TOBACCO ADVERTISING & PROMOTION					
Students who noticed anti-tobacco messages in the media^{21}	43.4	46.4	40.3*		
Students who noticed anti-tobacco messages at sporting or community events 27	39.1	43.2	35.0		
Current tobacco smokers who thought about quitting because of a warning label ²⁸	21.5	21.6	21.6		
Students who were taught in school about the dangers of tobacco use in the past 12 months	37.4	38.6	36.1		
KNOWLEDGE & ATTITUDES					
	OVERALL (%)	BOYS (%)	GIRLS (%)		
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	15.0	13.1	17.0*		
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	25.7	28.0	23.3		
Students who definitely thought other people's tobacco smoking is harmful to them	50.3	47.7	53.0		
Students who favored prohibiting smoking inside enclosed public places	83.8	80.6	87.1*		
Students who favored prohibiting smoking at outdoor public places	75.0	74.4	75.8		

¹ Smoked cigarettes, smoked other types of tobacco including shisha, used heated tobacco products, and/or used smokeless tobacco anytime during the past 30 days. ² Ever smoked tobacco including shisha, used heated tobacco products, and/or used smokeless tobacco. ³ Smoked cigarettes, shisha, or other types of tobacco anytime during the past 30 days. ⁴ Smoked cigarettes anytime during the past 30 days. ⁵ Smoked cigarettes on 20 or more days of the past 30 days. ⁶ Smoked tobacco other than cigarettes including shisha anytime during the past 30 days. ⁷ Smoked shisha anytime during the past 30 days. ⁸ Ever smoked cigarettes or other types of tobacco including shisha, even one or two puffs. ¹⁹ Ever smoked cigarettes, even one or two puffs. ¹⁹ Ever smoked tobacco other than cigarettes including shisha, even one or two puffs. ¹⁹ Ever smoked shisha, even one or two puffs. ¹⁹ Ever smoked shisha, even one or two puffs. ¹⁹ Ever smoked shisha, even one or two puffs. ¹⁹ Ever smoked tobacco other than cigarettes including shisha, even one or two puffs. ¹⁹ Ever smoked shisha, even one or two puffs. ¹⁹ Ever smoked tobacco anytime during the past 30 days. ¹⁹ Ever used heated tobacco products. ¹⁴ Used smokeless tobacco anytime during the past 30 days. ¹⁵ Ever used smokeles sbocco. ¹⁶ Susceptible to future tobacco use includes those who answered "definitely yes", "probably yes", or "probably not" to using tobacco if one of their best friends offered it to them, or "definitely yes", "probably yes", or "probably not" to using tobacco during the next 12 months. ¹⁷ Those who answered "Agree" or "Strongly Agree" to the statement: "I think I might enjoy smoking a cigarette". ¹⁸ Used electronic cigarettes anytime during the past 30 days. Current use of electronic cigarettes is assessed separately from cigarettes anytime during the past 30 days. ²⁰ During the past 30 days. ²

NOTE: Data are weighted to be nationally representative of all students aged 13-15 years who are enrolled in school. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.

⁻⁻Estimates based on unweighted cases less than 35 are not presented.

^{*}Gender comparisons are statistically significant at p < 0.05.