

# Slovakia (Ages 13-15)

## Global Youth Tobacco Survey (GYTS)

### FACT SHEET . . . . .



The Slovakia GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Slovakia could include in a comprehensive tobacco control program.

The Slovakia GYTS was a school-based survey of students in Forms 7 through 9 conducted in 2007. A two-stage cluster

sample design was used to produce representative data for all of Slovakia plus all 8 regions. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the class response rate was 99.6%, the student response rate was 86.1%, and the overall response rate was 85.7%. A total of 4,176 students aged 13-15 participated in the Slovakia GYTS.

### Prevalence

- 64.4% of students had ever smoked cigarettes (Boys = 68.7%, Girls = 60.2%)
- 26.6% currently use any tobacco product (Boys = 28.5%, Girls = 24.5%)
- 25.0% currently smoke cigarettes (Boys = 26.5%, Girls = 23.4%)
- 7.6% currently daily cigarette smokers (Boys = 9.3%, Girls = 5.8%)
- 10.2% currently smoke cigars (Boys = 11.3%, Girls = 8.9%)
- 28.8% ever smokers initiated smoking before age ten (Boys = 34.9%, Girls = 21.8%)
- 24.5% never smokers are likely to initiate smoking next year (Boys = 17.7%, Girls = 29.5%)

### Access and Availability - Current Smokers

- 5.8% usually smoke at home
- 51.4% buy cigarettes in a store
- 85.0% who bought cigarettes in a store were NOT refused purchase because of their age

### Exposure to Secondhand Smoke (SHS)

- 44.9% live in homes where others smoke in their presence
- 69.3% are around others who smoke in places outside their home
- 80.7% think smoking should be banned from public places
- 76.3% think smoke from others is harmful to them
- 49.9% have one or more parents who smoke
- 23.5% have most or all friends who smoke

### Cessation - Current Smokers

- 64.8% want to stop smoking
- 78.2% tried to stop smoking during the past year
- 65.9% have ever received help to stop smoking
- 16.6% always have or feel like having a cigarette first thing in the morning

### Media and Advertising

- 71.9% saw anti-smoking media messages vs. 66.4% saw pro media messages on TV
- 47.1% saw anti messages
- 53.0% saw anti smoking ads vs. 67.3% saw pro-cigarette ads in newspapers or magazines
- 17.3% have an object with a cigarette brand logo
- 5.5% were offered free cigarettes by a tobacco company representative

### School

- 58.1% had been taught in class, during the past year, about the dangers of smoking
- 48.9% had discussed in class, during the past year, reasons why people their age smoke
- 49.4% had been taught in class, during the past year, the effects of tobacco use

### Highlights

- Over one-quarter of the students currently use any form of tobacco; 1 in 4 currently smoke cigarettes; 7.6% are current daily smokers; 10.2% currently smoke cigars; 24.5% of never smokers are likely to initiate smoking next year.
- SHS exposure is very high – over 4 in 10 students live in homes where others smoke in their presence; 7 in 10 are exposed to smoke in public places; Half have one or more parents who smoke.
- Three-quarters of the students think smoke from others is harmful to them.
- Eight in 10 students think smoking in public places should be banned.
- Almost two-thirds of smokers want to quit.
- More than 1 in 6 students has an object with a cigarette brand logo on it.
- Seven in 10 students saw anti-smoking messages on TV vs. two-thirds saw pro messages; over half saw anti ads in newspapers or magazines vs. two-thirds saw pro ads.