Comoros (Ages 13-15) Global Youth Tobacco Survey (GYTS)



FACT SHEET

The Comoros GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Comoros could include in a comprehensive tobacco control program.

The Comoros GYTS was a school-based survey of students in 6ème, 5ème, 4ème, and 3ème conducted in 2007.

A two-stage cluster sample design was used to produce representative data for all of Comoros. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, the class response rate was 100%, the student response rate was 80.8%, and the overall response rate was 80.8%. A total of 811 students aged 13-15 participated in the Comoros GYTS.

Prevalence

26.5% of students had ever smoked cigarettes (Boy = 39.3%, Girl = 17.2%)

18.1% currently use any tobacco product (Boy = 21.8%, Girl = 14.8%)

9.6% currently smoke cigarettes (Boy = 13.5%, Girl = 6.9%)

11.4% currently use other tobacco products (Boy = 12.5%, Girl = 9.9%)

9.9% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

46.9% think boys and 22.4% think girls who smoke have more friends 20.2% think boys and 16.9% think girls who smoke look more attractive

Access and Availability - Current Smokers

25.9% usually smoke at home 23.4% buy cigarettes in a store

Exposure to Secondhand Smoke (SHS)

35.2% live in homes where others smoke in their presence

58.3% are around others who smoke in places outside their home

78.8% think smoking should be banned from public places

62.9% think smoke from others is harmful to them

23.7% have one or more parents who smoke

6.9% have most or all friends who smoke

Cessation - Current Smokers

100.0% have ever received help to stop smoking

Media and Advertising

77.6% saw anti-smoking media messages, in the past 30 days

51.5% saw pro-cigarette ads on billboards, in the past 30 days

48.3% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

20.1% have an object with a cigarette brand logo

6.4% were offered free cigarettes by a tobacco company representative

School

36.5% had been taught in class, during the past year, about the dangers of smoking 24.8% had discussed in class, during the past year, reasons why people their age smoke 30.4% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 18.1% of students currently use any form of tobacco; 9.6% currently smoke cigarettes; 11.4% currently use some other form of tobacco.
- SHS exposure is high More than one-third of the students live in homes where others smoke in their presence, and over half are exposed to smoke in public places; Nearly one-quarter have at least one parent who smokes, and 6.9% have most or all friends who smoke.
- Nearly two-thirds of the students think smoke from others is harmful to them.
- Over three-quarters of the students think smoking in public places should be banned.
- One in 5 students has an object with a cigarette brand logo on it.
- Nearly 8 in 10 students saw antismoking media messages in the past 30 days; Half the students saw pro-cigarette ads on billboards and in newspapers or magazines in the past 30 days.