

The Ecuador-Quito GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Ecuador-Quito could include in a comprehensive tobacco control program.

The Ecuador-Quito GYTS was a school-based survey of students in 8 vo de basica, 9 no de basica, 10 mo de basica and 1ro de bachillerato conducted in 2007.

A two-stage cluster sample design was used to produce representative data for all Ecuador-Quito. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 92.0%, the class response rate was 100.0%, the student response rate was 95.1% and the overall response rate was 87.5%. A total of 1,389 students aged 13-15 participated in the Ecuador-Quito GYTS.

Prevalence

56.0% of students had ever smoked cigarettes (Boys = 62.6%, Girls = 50.2%) 28.6% currently use any tobacco product (Boys = 31.2%, Girls = 26.1%) 20.5% currently smoke cigarettes (Boys = 23.2%, Girls = 18.1%) 15.3% currently use other tobacco products (Boys = 15.9%, Girls = 14.6%) 28.0% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

20.9% think boys and 30.6% think girls who smoke have more friends 13.0% think boys and 9.8% think girls who smoke look more attractive

Access and Availability - Current Smokers

16.0% usually smoke at home41.4% buy cigarettes in a store58.2% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

28.9% live in homes where others smoke in their presence

52.5% are around others who smoke in places outside their home

86.9% think smoking should be banned from public places

75.5% think smoke from others is harmful to them

39.1% have one or more parents who smoke

13.0% have most or all friends who smoke

Cessation - Current Smokers

59.2% want to stop smoking62.4% tried to stop smoking during the past year75.3% have ever received help to stop smoking

Media and Advertising

82.1% saw anti-smoking media messages, in the past 30 days

82.3% saw pro-cigarette ads on billboards, in the past 30 days

74.1% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

12.6% have an object with a cigarette brand logo

15.8% were offered free cigarettes by a tobacco company representative

School

53.1% had been taught in class, during the past year, about the dangers of smoking 21.8% had discussed in class, during the past year, reasons why people their age smoke 45.2% had been taught in class, during the past year, the effects of tobacco use

Highlights

- One in 5 students currently smokes cigarettes; almost 3 in 10 students use any tobacco products; and 28.0% of never smokers are likely to initiate smoking next year.
- SHS exposure is high almost 3 in 10 students live in homes where others smoke; over half the students are exposed to smoke of others outside their home; 2 in 5 have at least one parent who smokes.
- Three-quarters of the students think smoke from others is harmful to them.
- Over 8 in 10 students think smoking in public places should be banned.
- Three in 5 current smokers want to stop smoking.
- One in 7 students was offered a free cigarette by a tobacco company representative.
- Over 8 in 10 students saw antismoking media messages in the past 30 days; more than 8 in 10 students saw pro-cigarette ads on billboards and three-quarters of the students saw pro-cigarette ads in newspapers or magazines in the past 30 days.