Estonia (Ages 13-15)	youth
Global Youth Tobacco Survey (GYTS)	tobacco
FACT SHEET · · · · · · · · · · · ·	survey

The Estonia GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Estonia could include in a comprehensive tobacco control program.

The Estonia GYTS was a school-based survey of students in grades 7 through 9 conducted in 2003. A two-stage cluster

sample design was used to produce representative data for all of Estonia. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 95.7%, the class response rate was 100%, the student response rate was 81.6%, and the overall response rate was 78.2%. A total of 4,307 students aged 13-15 participated in the Estonia GYTS.

Prevalence

76.0% of students had ever smoked cigarettes (Boy = 80.7%, Girl = 72.0%) 29.5% currently use any tobacco product (Boy = 31.2%, Girl = 27.3%) 28.9% currently smoke cigarettes (Boy = 29.8%, Girl = 27.4%)

10.9% currently daily cigarette smokers (Boy = 12.9%, Girl = 9.0%)

12.6% currently smoke cigars (Boy = 12.9%, Girl = 11.6%)

36.6% ever smokers initiated smoking before age ten (Boy = 47.0%, Girl = 27.2%)

36.6% of never smokers are likely to initiate smoking next year (Boy = 33.1%, Girl = 38.3%)

Access and Availability - Current Smokers

7.2% usually smoke at home

41.2% buy cigarettes in a store

64.6% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

80.6% live in homes where others smoke in their presence
90.7% are around others who smoke in places outside their home
78.3% think smoking should be banned from public places
57.6% think smoke from others is harmful to them
58.7% have one or more parents who smoke
27.6% have most or all friends who smoke

Cessation - Current Smokers

61.0% want to stop smoking70.5% tried to stop smoking during the past year93.4% have ever received help to stop smoking

15.4% always have or feel like having a cigarette first thing in the morning

Media and Advertising

52.5% saw anti-smoking media messages vs. 75.8% saw pro media messages on TV
64.2% saw anti messages vs. 60.9% saw pro messages on billboards
39.3% saw anti smoking ads vs. 60.4% saw pro-cigarette ads in newspapers or magazines
26.3% have an object with a cigarette brand logo

15.2% were offered free cigarettes by a tobacco company representative

School

59.0% had been taught in class, during the past year, about the dangers of smoking 45.3% had discussed in class, during the past year, reasons why people their age smoke 47.6% had been taught in class, during the past year, the effects of tobacco use

Highlights

- One in 3 students currently use any form of tobacco; 28.9% currently smoke cigarettes; 10.9% current daily smokers; 12.6% currently smoke cigars; Over one-third of never smokers likely to initiate smoking next year.
- SHS exposure is very high 8 in 10 students live in homes where others smoke in their presence; 9 in 10 are exposed to smoke in public places; Almost 6 in 10 have one or more parents who smoke.
- Almost 3 in 5 students think smoke from others is harmful to them.
- More than three-quarters of the students think smoking in public places should be banned.
- Over 3 in 5 smokers want to quit.
- More than one-quarter of the students has an object with a cigarette brand logo.
- Over half the students saw antismoking messages on TV vs. three-quarters saw pro messages; Close to two-thirds saw anti messages on billboards vs. 6 in 10 saw pro messages; 4 in 10 saw anti ads in newspapers vs. 6 in 10 saw pro ads.