

The Mauritania GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Mauritania could include in a comprehensive tobacco control program.

The Mauritania GYTS was a school-based survey of students in 1ère AS, 2 ème AS, 3 ème AS, and 4 ème AS conducted in 2009.

A two-stage cluster sample design was used to produce representative data for Mauritania. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the class response rate was 100.0%, the student response rate was 91.1%, and the overall response rate was 91.1%. A total of 1,769, students aged 13-15 participated in the Mauritania GYTS.

Prevalence

22.1% of students had ever smoked cigarettes (Boy = 28.5%, Girl = 16.1%) 22.6% currently use any tobacco product (Boy = 27.5%, Girl = 17.7%) 11.6% currently smoke cigarettes (Boy = 14.6%, Girl = 9.0%) 13.1% currently use other tobacco product (Boy = 15.0%, Girl = 10.2%)

13.1% currently use other tobacco products (Boy = 15.9%, Girl = 10.2%)

19.7% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

36.0% think boys and 27.8% think girls who smoke have more friends 31.0% think boys and 23.4% think girls who smoke look more attractive

Access and Availability - Current Smokers

12.1% usually smoke at home

49.4% buy cigarettes in a store

78.5% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

37.5% live in homes where others smoke in their presence

- $50.9\%\,$ are around others who smoke in places outside their home
- $58.7\%\,$ think smoking should be banned from public places
- 38.9% think smoke from others is harmful to them

23.0% have one or more parents who smoke

8.8% have most or all friends who smoke

Cessation - Current Smokers

50.9% want to stop smoking61.2% tried to stop smoking during the past year

Media and Advertising

74.1% saw anti-smoking media messages, in the past 30 days

- 61.8% saw pro-cigarette ads on billboards, in the past 30 days
- 53.8% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 21.6% have an object with a cigarette brand logo

School

43.1% had been taught in class, during the past year, about the dangers of smoking 31.9% had discussed in class, during the past year, reasons why people their age smoke 37.6% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 22.6% of students currently use any form of tobacco; 11.6% currently smoke cigarettes; 13.1% currently use some other form of tobacco.
- SHS exposure is moderate more than one-third of the students live in homes where others smoke, and about half of the students are exposed to smoke around others outside of the home; Nearly one-quarter of the students have at least one parent who smokes.
- Nearly two in five students think smoke from others is harmful to them.
- Almost three in five students think smoking in public places should be banned.
- Half of the current smokers want to stop smoking.
- More than one in five students have an object with a cigarette brand logo on it.
- Three-quarters of the students saw anti-smoking media messages in the past 30 days; three in five students saw procigarette ads on billboards and over half saw pro-tobacco ads in newspapers or magazines in the past 30 days.