Syria UNRWA (Ages 13-15)	global
Global Youth Tobacco Survey (GYTS)	youth tobacco
FACT SHEET · · · · · · · · · · · ·	survey

The Syria UNRWA GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Syria UNRWA could include in a comprehensive tobacco control program.

The Syria UNRWA GYTS was a school-based survey of students in grades 7, 8, and 9 conducted in 2008. A two-stage cluster sample

design was used to produce representative data for all of Syria UNRWA. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100%, class response rate was 100%, the student response rate was 95.9%, and the overall response rate was 95.9%. A total of 1,390 students aged 13-15 participated in the Syria UNRWA GYTS.

Prevalence

29.6% of students had ever smoked cigarettes (Boy = 39.5%, Girl = 18.5%) 42.4% currently use any tobacco product (Boy = 49.6%, Girl = 33.8%) 13.3% currently smoke cigarettes (Boy = 19.6%, Girl = 6.3%) 31.2% currently smoke shsisha (Boy = 36.1%, Girl = 24.8%) 37.7% currently use other tobacco products (Boy = 42.7%, Girl = 31.3%) 20.7% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

28.1% think boys and 21.2% think girls who smoke have more friends 21.8% think boys and 18.4% think girls who smoke look more attractive

Access and Availability - Current Smokers

25.0% usually smoke at home 35.5% buy cigarettes in a store 82.4% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

66.1% live in homes where others smoke in their presence 67.6% are around others who smoke in places outside their home 80.1% think smoking should be banned from public places 67.7% think smoke from others is harmful to them 57.0% have one or more parents who smoke 13.9% have most or all friends who smoke

Cessation - Current Smokers

73.1% want to stop smoking 71.3% tried to stop smoking during the past year 75.2% have ever received help to stop smoking

Media and Advertising

70.4% saw anti-smoking media messages, in the past 30 days

72.3% saw pro-cigarette ads on billboards, in the past 30 days

59.4% saw pro-cigarette ads in newspapers or magazines, in the past 30 days

13.9% have an object with a cigarette brand logo

13.6% were offered free cigarettes by a tobacco company representative

School

67.5% had been taught in class, during the past year, about the dangers of smoking 46.8% had discussed in class, during the past year, reasons why people their age smoke 54.5% had been taught in class, during the past year, the effects of tobacco use

Highlights

- More than 2 in 5 students currently use any form of tobacco; 13.3% currently smoke cigarettes; 37.7% currently use other tobacco products.
- Over 3 in 10 students currently smoke shsisha
- SHS exposure is very high nearly two-thirds of students live in homes where others smoke; 67.6% of students are exposed to smoke in public places; Almost 3 in 5 students have one or more parents who smoke.
- Two-thirds of students think that smoke from others is harmful to them.
- Four in 5 students think smoking should be banned in public places.
- Almost three-quarters of current smokers want to stop.
- 13.9% of students have an object with a cigarette brand logo on it.
- Seven in 10 students saw antismoking media messages in the past 30 days; close to threequarters of the students saw procigarette ads on billboards in the past 30 days and 3 in 5 students saw pro-cigarette ads in newspapers or magazines.