GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:



Monitor tobacco use & prevention policies

Protect people from tobacco smoke

Offer help to quit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising, promotion, & sponsorship

Raise taxes on tobacco

GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco media and advertising, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is self-administered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In Trinidad and Tobago, GYTS was conducted in 2017 by the Ministry of Health. The overall response rate was 90.7%. A total of 4,128 eligible students in forms 1 - 5, 6 lower and 6 upper completed the survey, of which 2,292 were aged 13-15 years. Data are reported for students aged 13-15 years.

GYTS Highlights

TOBACCO USE

- 14.0% of students, 17.3% of boys, and 10.8% of girls currently used any tobacco products.
- 11.0% of students, 13.6% of boys, and 8.6% of girls currently smoked tobacco.
- 6.7% of students, 8.6% of boys, and 4.9% of girls currently smoked cigarettes.
- 4.1% of students, 5.0% of boys, and 3.2% of girls currently used smokeless tobacco.

ELECTRONIC CIGARETTES

• 17.2% of students, 21.7% of boys, and 12.9% of girls currently used electronic cigarettes.

CESSATION

- Almost 6 in 10 (56.8%) current tobacco smokers tried to stop smoking in the past 12 months.
- More than 4 in 10 (43.6%) current tobacco smokers wanted to stop smoking now.

SECONDHAND SMOKE

- 25.8% of students were exposed to tobacco smoke at home.
- 30.2% of students were exposed to tobacco smoke inside enclosed public places.

ACCESS & AVAILABILITY

- 42.8% of current cigarette smokers bought cigarettes from a store, shop, street vendor, or kiosk.
- Among current cigarette smokers who tried to buy cigarettes, 62.1% were not prevented from buying them because of their age.

MEDIA

- More than 4 in 10 (44.1%) students noticed anti-tobacco messages in the media.
- Almost 3 in 10 (28.7%) students noticed tobacco advertisements or promotions when visiting points of sale.
- More than 1 in 10 (14.1%) students had something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 53.2% of students definitely thought other people's tobacco smoking is harmful to them.
- 77.5% of students favored prohibiting smoking inside enclosed public places.










| TOBACCO USE | | | |
|--|-------------|----------|-----------|
| SMOKED TOBACCO | OVERALL (%) | BOYS (%) | GIRLS (%) |
| Current tobacco smokers ¹ | 11.0 | 13.6 | 8.6* |
| Current cigarette smokers ² | 6.7 | 8.6 | 4.9* |
| Frequent cigarette smokers ³ | 1.9 | 3.1 | 0.8* |
| Current smokers of other tobacco ⁴ | 6.2 | 7.4 | 4.9* |
| Ever tobacco smokers ⁵ | 31.5 | 35.7 | 27.5* |
| Ever cigarette smokers ⁶ | 25.2 | 30.3 | 20.5* |
| Ever smokers of other tobacco ⁷ | 13.9 | 15.2 | 12.7 |
| SMOKELESS TOBACCO | | | |
| Current smokeless tobacco users ⁸ | 4.1 | 5.0 | 3.2* |
| Ever smokeless tobacco users ⁹ | 8.5 | 9.8 | 7.2 |
| ANY TOBACCO USE (smoked and/or smokeless) | | | |
| Current tobacco users ¹⁰ | 14.0 | 17.3 | 10.8* |
| Ever tobacco users ¹¹ | 35.5 | 40.3 | 30.9* |
| SUSCEPTIBILITY | | | |
| Never to bacco users susceptible to to bacco use in the ${\rm future^{12}}$ | 17.1 | 18.8 | 15.7 |
| Never to bacco smokers who thought they might enjoy smoking a cigarette $^{\! 13}$ | 7.8 | 8.0 | 7.7 |



ELECTRONIC CIGARETTES

| | OVERALL (%) | BOYS (%) | GIRLS (%) |
|--|-------------|----------|-----------|
| Current electronic cigarette users ¹⁴ | 17.2 | 21.7 | 12.9* |
| Ever electronic cigarette users ¹⁵ | 27.7 | 32.6 | 23.1* |

CESSATION

| | OVERALL (%) | BOYS (%) | GIRLS (%) |
|---|-------------|----------|-----------|
| Current tobacco smokers who tried to stop smoking in the past 12 months | 56.8 | 53.9 | 62.4 |
| Current tobacco smokers who wanted to stop smoking now | 43.6 | 44.1 | 42.7 |
| Current tobacco smokers who thought they would be able to stop smoking if they wanted to | 78.8 | 78.4 | 79.4 |
| Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking | 22.8 | 25.4 | 18.1 |

SECONDHAND SMOKE

| | OVERALL (%) | BOYS (%) | GIRLS (%) |
|--|-------------|----------|-----------|
| Exposure to tobacco smoke at home ¹⁶ | 25.8 | 24.6 | 26.9 |
| Exposure to tobacco smoke inside any enclosed public place 16 | 30.2 | 29.5 | 31.0 |
| Exposure to tobacco smoke at any outdoor public place ¹⁶ | 37.2 | 35.7 | 38.6 |
| Students who saw anyone smoking inside the school building or outside on school property ¹⁷ | 31.9 | 33.1 | 30.8 |

| ACCESS & AVAILABILITY | | | |
|---|-------------|----------|-----------|
| | OVERALL (%) | BOYS (%) | GIRLS (%) |
| Current cigarette smokers who bought cigarettes from a store, shop, street vendor, or kiosk ¹⁸ | 42.8 | 54.9 | 20.3* |
| Current cigarette smokers who were not prevented from buying cigarettes because of their age^{19} | 62.1 | 68.0 | |
| Current cigarette smokers who bought cigarettes as individual sticks ²⁰ | 31.3 | 28.4 | |

| MEDIA | | | |
|---|-------------|----------|-----------|
| TOBACCO ADVERTISING | OVERALL (%) | BOYS (%) | GIRLS (%) |
| Students who noticed to bacco advertisements or promotions at points of sale 21 | 28.7 | 30.8 | 26.6 |
| Students who saw anyone using tobacco on television, videos, or movies ²² | 64.3 | 63.0 | 65.6 |
| Students who were ever offered a free tobacco product from a tobacco company representative | 8.6 | 10.5 | 6.8* |
| Students who had something with a tobacco brand logo on it | 14.1 | 14.8 | 13.3 |
| ANTI-TOBACCO ADVERTISING | | | |
| Students who noticed anti-tobacco messages in the media^{17} | 44.1 | 41.7 | 46.2 |
| Students who noticed anti-tobacco messages at sporting or community events 23 | 33.7 | 37.9 | 28.9* |
| Current tobacco smokers who thought about quitting because of a warning label ²⁴ | 30.5 | 32.3 | 27.1 |
| Students who were taught in school about the dangers of tobacco use in the past 12 months | 38.5 | 41.2 | 36.0 |

KNOWLEDGE & ATTITUDES

| | OVERALL (%) | BOYS (%) | GIRLS (%) |
|---|-------------|----------|-----------|
| Students who definitely thought it is difficult to quit once someone starts smoking tobacco | 30.7 | 26.9 | 34.3* |
| Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings | 32.0 | 35.4 | 28.8* |
| Students who definitely thought other people's tobacco smoking is harmful to them | 53.2 | 50.4 | 55.8* |
| Students who favored prohibiting smoking inside enclosed public places | 77.5 | 71.5 | 83.1* |
| Students who favored prohibiting smoking at outdoor public places | 62.3 | 58.3 | 66.1* |

¹ Smoked cigarettes or other type of tobacco anytime during the past 30 days. ² Smoked cigarettes anytime during the past 30 days. ³ Smoked cigarettes on 20 or more days of the past 30 days. ⁴ Smoked tobacco other than cigarettes anytime during the past 30 days. ⁵ Ever smoked cigarettes or other type of tobacco, even one or two puffs. ⁶ Ever smoked cigarettes, even one or two puffs. ⁶ Ever smoked cigarettes, even one or two puffs. ⁷ Ever smoked tobacco other than cigarettes, even one or two puffs. ⁸ Used smokeless tobacco anytime during the past 30 days. ⁹ Ever used smokeless tobacco. ¹⁰ Smoked cigarettes, smoked other type of tobacco, and/or used smokeless tobacco anytime during the past 30 days. ¹¹ Ever smoked tobacco and/or used smokeless tobacco. ¹² Susceptible to future tobacco use includes those who answered "definitely yes", "probably yes", or "probably not" to using tobacco if one of their best friends offered it to them, or "definitely yes", "probably yes", or "probably not" to using tobacco during the next 12 months. ¹³ Those who answered "Agree" or "Strongly Agree" to the statement: "I think I might enjoy smoking a cigarette". ¹⁴ Used electronic cigarettes anytime during the past 30 days. Current use of electronic cigarettes is assessed separately from cigarettes, other smoked tobacco products, and smokeless tobacco products and is not included in the current definition of current any tobacco use. ¹⁵ Ever used electronic cigarettes in their entire life. ¹⁶ During the past 7 days. ¹⁷ During the past 30 days. ¹⁹ Among those who tried to buy cigarettes during the past 30 days. ²⁰ Based on the last purchase, among those who bought cigarettes during the past 30 days. ²¹ Among those who visited a point of sale in the past 30 days. ²² Among those who watched television, videos, or movies in the past 30 days. ²³ Among those who attended sporting or community events in the past 30 days. ²⁴ Among those who noticed warning labels on cigarette packages in the past

NOTE: Data are weighted to be nationally representative of all students aged 13-15 years who are enrolled in school. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.

⁻⁻Estimates based on unweighted cases less than 35 are not presented.

^{*}Gender comparisons are significant at p < 0.05.