GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:

> Monitor tobacco use & prevention policies

Protect people from tobacco smoke

Offer help to quit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising, promotion, & sponsorship

Raise taxes on tobacco

GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco media and advertising, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is selfadministered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In Uganda, GYTS was conducted in 2018 by the Ministry of Health. The overall response rate was 75.8%. A total of 3,458 eligible students in primary 7 and senior 1-3 of secondary school completed the survey, of which 2,068 were aged 13-15 years. Data are reported for students aged 13-15 years.

GYTS Highlights

TOBACCO USE

- 10.5% of students, 11.7% of boys, and 9.4% of girls currently used any tobacco products.
- 5.5% of students, 6.8% of boys, and 4.3% of girls currently smoked tobacco.
- 3.5% of students, 4.7% of boys, and 2.4% of girls currently smoked cigarettes.
- 6.5% of students, 7.1% of boys, and 6.0% of girls currently used smokeless tobacco.

CESSATION

• More than 9 in 10 (93.0%) current tobacco smokers tried to stop smoking in the past 12 months.

SECONDHAND SMOKE

- 19.2% of students were exposed to tobacco smoke at home.
- 30.0% of students were exposed to tobacco smoke inside enclosed public places.

ACCESS & AVAILABILITY

• 22.9% of current cigarette smokers bought cigarettes from a supermarket, shop, street vendor, or kiosk.

MEDIA

- Almost 6 in 10 (58.2%) students noticed anti-tobacco messages in the media.
- Almost 4 in 10 (39.3%) students noticed tobacco advertisements or promotions when visiting points of sale.
- Almost 1 in 10 (8.5%) students had something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 56.9% of students definitely thought other people's tobacco smoking is harmful to them.
- 68.6% of students favored prohibiting smoking inside enclosed public places.








The mark "CDC" is owned by the US Dept. of Health and Human Services and is used with permission. Use of this logo is not an endorsement by HHS or CDC of any particular product, service, or enterprise.

GYTS GLOBAL YOUTH TOBACCO SURVEY

FACT SHEET UGANDA 2018

TOBACCO USE

SMOKED TOBACCO	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers ¹	5.5	6.8	4.3
Current cigarette smokers ²	3.5	4.7	2.4
Frequent cigarette smokers ³	0.3	0.3	0.2
Current smokers of other tobacco ⁴	3.0	3.1	2.9
Ever tobacco smokers ⁵	21.3	28.5	14.3*
Ever cigarette smokers ⁶	16.0	23.4	8.7*
Ever smokers of other tobacco ⁷	11.4	13.7	9.4*
SMOKELESS TOBACCO			
Current smokeless tobacco users ⁸	6.5	7.1	6.0
Ever smokeless tobacco users ⁹	11.8	12.6	11.0
ANY TOBACCO USE (smoked and/or smokeless)			
Current tobacco users ¹⁰	10.5	11.7	9.4
Ever tobacco users ¹¹	27.8	33.7	22.1*
SUSCEPTIBILITY			
Never to bacco users susceptible to to bacco use in the $\rm future^{12}$	34.6	33.1	35.6
Never tobacco smokers who thought they might enjoy smoking a cigarette ¹³	8.1	8.9	7.4



CESSATION

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers who tried to stop smoking in the past 12 months	93.0		
Current tobacco smokers who thought they would be able to stop smoking if they wanted to	88.7		
Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking	30.0	25.4	

SECONDHAND SMOKE

	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home ¹⁴	19.2	19.6	19.0
Exposure to tobacco smoke inside any enclosed public place ¹⁴	30.0	34.6	25.9
Exposure to tobacco smoke at any outdoor public place ¹⁴	32.4	34.3	31.1
Students who saw anyone smoking inside the school building or outside on school property ¹⁵	28.8	26.6	31.1

ACCESS & AVAILABILITY

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current cigarette smokers who bought cigarettes from a supermarket, shop, street vendor, or kiosk ¹⁶	22.9		
MEDIA			
TOBACCO ADVERTISING	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who noticed tobacco advertisements or promotions at points of sale ¹⁷	39.3	36.7	42.0
Students who saw anyone using tobacco on television, videos, or movies ¹⁸	64.9	62.5	67.7
Students who were ever offered a free tobacco product from a tobacco company representative	6.9	6.3	7.1
Students who had something with a tobacco brand logo on it	8.5	8.6	8.0
ANTI-TOBACCO ADVERTISING			
Students who noticed anti-tobacco messages in the media ¹⁵	58.2	60.2	56.3
Students who noticed anti-tobacco messages at sporting or community events ¹⁹	47.4	50.8	43.7
Current tobacco smokers who thought about quitting because of a warning label ²⁰	39.5	43.1	34.4
Students who were taught in school about the dangers of tobacco use in the past 12 months	76.9	76.0	78.2

KNOWLEDGE & ATTITUDES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	24.7	26.7	23.1
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	15.8	17.2	14.5
Students who definitely thought other people's tobacco smoking is harmful to them	56.9	58.7	55.1
Students who favored prohibiting smoking inside enclosed public places	68.6	69.2	68.4
Students who favored prohibiting smoking at outdoor public places	65.6	66.2	65.1

¹ Smoked cigarettes or other type of tobacco anytime during the past 30 days. ² Smoked cigarettes anytime during the past 30 days. ³ Smoked cigarettes on 20 or more days of the past 30 days. ⁴ Smoked tobacco other than cigarettes anytime during the past 30 days. ⁵ Ever smoked cigarettes or other type of tobacco, even one or two puffs. ⁶ Ever smoked cigarettes, even one or two puffs. ⁶ Ever smoked cigarettes, smoked or used smokeless tobacco. ¹⁰ Smoked cigarettes, smoked other type of tobacco, and/or used smokeless tobacco. ¹² Susceptible to future tobacco anytime during the past 30 days. ¹¹ Ever smoked definitely yes", "probably yes", or "probably not" to using tobacco diring the next 12 months. ¹³ Those who answered "definitely yes", "probably yes", or "probably not" to using tobacco during the next 12 months. ¹³ Those who answered "affinitely ass. ¹⁸ During the past 30 days. ¹⁷ Among those who attended sporting or community events in the past 30 days. ¹⁰ Among those who attended sporting or community events in the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who attended the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who attended the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who attended the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who stended the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who attended the past 30 days. ²⁰ Among those who attended the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in th

NOTE: Data are weighted to be nationally representative of all students aged 13-15 years who are enrolled in school. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.

--Estimates based on unweighted cases less than 35 are not presented. *Gender comparisons are significant at p < 0.05.