## **GYTS Objectives**

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:



Monitor tobacco use & prevention policies

Protect people from tobacco smoke

Offer help to quit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising, promotion, & sponsorship

Raise taxes on tobacco

# **GYTS Methodology**

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco media and advertising, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is self-administered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In Uruguay, GYTS was conducted in 2014 by the National Drug Board, under the coordination of the – Ministerio de Salud Publica. The overall response rate was 85.9%. A total of 4,902 eligible students in 1<sup>st</sup>-3<sup>rd</sup> Basic Cycle and 4<sup>th</sup> year (1<sup>st</sup> year bacharelate) completed the survey, of which 3,256 were aged 13-15 years. Data are reported for students aged 13-15 years.

## **GYTS Highlights**

#### TOBACCO USE

- 12.8% of students, 12.7% of boys, and 12.5% of girls currently used any tobacco products.
- 9.9% of students, 9.6% of boys, and 9.8% of girls currently smoked tobacco.
- 8.2% of students, 7.2% of boys, and 8.7% of girls currently smoked cigarettes.
- 3.5% of students, 4.0% of boys, and 3.1% of girls currently used smokeless tobacco.
- 67.6% of students, 59.7% of boys, and 71.7% of girls who are current smokers showed signs of smoking dependence.

#### **CESSATION**

- More than 5 in 10 current smokers tried to stop smoking in the past 12 months.
- More than 3 in 10 current smokers wanted to stop smoking now.

#### **SECONDHAND SMOKE**

- 40.6% of students were exposed to tobacco smoke at home.
- 40.1% of students were exposed to tobacco smoke inside enclosed public places.

#### **ACCESS & AVAILABILITY**

- 64.7% of current cigarette smokers bought cigarettes from a store, shop, street vendor, kiosk, parlor, newsstand, supermarket, gas station, tavern, bar or restaurant.
- Among current cigarette smokers who tried to buy cigarettes, 74.1% were not prevented from buying them because of their age.

### **MEDIA**

- Almost 6 in 10 students noticed anti-tobacco messages in the media.
- Almost 6 in 10 students noticed tobacco advertisements or promotions when visiting points of sale.
- Almost 1 in 10 students had something with a tobacco brand logo on it.

#### **KNOWLEDGE & ATTITUDES**

- 57.0% of students definitely thought other people's tobacco smoking is harmful to them.
- 86.6% of students favored prohibiting smoking inside enclosed public places.














TOBACCO USE			
SMOKED TOBACCO	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers <sup>1</sup>	9.9	9.6	9.8
Current cigarette smokers <sup>2</sup>	8.2	7.2	8.7
Frequent cigarette smokers <sup>3</sup>	1.6	1.5	1.8
Current smokers of other tobacco <sup>4</sup>	2.7	3.4	2.1
Ever tobacco smokers <sup>5</sup>	28.5	27.2	29.5
Ever cigarette smokers <sup>6</sup>	25.9	23.9	27.5*
Ever smokers of other tobacco <sup>7</sup>	7.3	8.4	6.2*
SMOKELESS TOBACCO			
Current smokeless tobacco users <sup>8</sup>	3.5	4.0	3.1
Ever smokeless tobacco users <sup>9</sup>	6.2	6.5	5.8
ANY TOBACCO USE (smoked and/or smokeless)			
Current tobacco users <sup>10</sup>	12.8	12.7	12.5
Ever tobacco users <sup>11</sup>	32.3	31.1	33.0
SUSCEPTIBILITY			
Never to bacco users susceptible to to bacco use in the future $^{\rm 12}$	16.6	14.1	18.5*
Never smokers who thought they might enjoy smoking a cigarette $^{13}$	7.8	9.3	6.6
SMOKING DEPENDENCY			
Current smokers who are showing signs of smoking dependence <sup>14</sup>	67.6	59.7	71.7



### **CESSATION**

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current smokers who tried to stop smoking in the past 12 months	51.2	48.5	52.5
Current smokers who wanted to stop smoking now	32.1	28.3	33.7
Current smokers who thought they would be able to stop smoking if they wanted to	84.9	79.0	88.3
Current smokers who have ever received help/advice from a program or professional to stop smoking	17.8	20.0	15.6

SECONDHAND SMOKE			
	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home <sup>15</sup>	40.6	35.5	44.8*
Exposure to tobacco smoke inside any enclosed public place <sup>15</sup>	40.1	34.0	45.1*
Exposure to tobacco smoke at any outdoor public place <sup>15</sup>	46.8	40.9	51.8*
Students who saw anyone smoking inside the school building or outside on school property <sup>16</sup>	37.0	35.7	38.1

ACCESS & AVAILABILITY			
	OVERALL (%)	BOYS (%)	GIRLS (%)
Current cigarette smokers who bought cigarettes from a store, shop, street vendor, kiosk, parlor, newsstand, supermarket, gas station, tavern, bar or restaurant <sup>17</sup>	64.7	68.9	62.2
Current cigarette smokers who were not prevented from buying cigarettes because of their age $^{18}$	74.1	71.4	77.5
Current cigarette smokers who bought cigarettes as individual sticks <sup>19</sup>	41.5	36.5	45.7
MEDIA			
TOBACCO ADVERTISING	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who noticed to bacco advertisements or promotions at points of sale $^{20}$	58.1	55.8	59.9*
Students who saw anyone using tobacco on television, videos, or movies <sup>21</sup>	75.4	74.8	75.9
Students who were ever offered a free tobacco product from a tobacco company representative	5.7	7.0	4.5*
Students who had something with a tobacco brand logo on it	7.5	9.0	5.8*
ANTI-TOBACCO ADVERTISING			
Students who noticed anti-tobacco messages in the media $^{16}$	59.9	56.2	63.2*
Students who noticed anti-tobacco messages at sporting or community events <sup>22</sup>	39.3	42.6	35.8*
Current smokers who thought about quitting because of a warning label <sup>23</sup>	33.1	25.6	37.5
Students who were taught in school about the dangers of tobacco use in the past 12 months	33.0	34.2	32.1
KNOWLEDGE & ATTITUDES	5		
	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	19.4	19.0	19.9
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	24.8	29.0	21.2*
Students who definitely thought other people's tobacco smoking is harmful to them	57.0	57.2	57.0
Students who favored prohibiting smoking inside enclosed public places	86.6	84.9	88.2*

<sup>1</sup> Smoked cigarettes or other type of tobacco anytime during the past 30 days. <sup>2</sup> Smoked cigarettes anytime during the past 30 days. <sup>3</sup> Smoked cigarettes on 20 or more days of the past 30 days. <sup>4</sup> Smoked tobacco other than cigarettes anytime during the past 30 days. <sup>5</sup> Ever smoked cigarettes or other type of tobacco, even one or two puffs. <sup>6</sup> Ever smoked cigarettes, even one or two puffs. <sup>7</sup> Ever smoked tobacco other than cigarettes, even one or two puffs. <sup>8</sup> Used smokeless tobacco anytime during the past 30 days. <sup>9</sup> Ever used smokeless tobacco. <sup>10</sup> Smoked cigarettes, smoked other type of tobacco, and/or used smokeless tobacco anytime during the past 30 days. <sup>11</sup> Ever smoked tobacco and/or used smokeless tobacco. <sup>12</sup> Susceptible to future tobacco use includes those who answered "definitely yes", "probably yes", or "probably not" to using tobacco if one of their best friends offered it to them, or "definitely yes", "probably yes", or "probably not" to using tobacco during the next 12 months. <sup>13</sup> Those who answered "Agree" or "Strongly Agree" to the statement: "I think I might enjoy smoking a cigarette". <sup>14</sup> Smoking dependency includes those who "sometimes" or "always" smoke or feel like smoking tobacco first thing in the morning, or those who start to feel a strong desire to smoke again within one full day after smoking tobacco. <sup>15</sup> During the past 7 days. <sup>16</sup> During the past 30 days. <sup>17</sup> Outlet from which current cigarette smokers bought cigarettes the last time they smoked cigarettes in the past 30 days. <sup>18</sup> Among those who tried to buy cigarettes during the past 30 days. <sup>19</sup> Based on the last purchase, among those who bought cigarettes during the past 30 days. <sup>20</sup> Among those who watched television, videos, or movies in the past 30 days. <sup>22</sup> Among those who watched television, videos, or movies in the past 30 days. <sup>23</sup> Among those who watched television, videos, or movies in the past 30 days. <sup>24</sup> Among those who watched television, videos, or movies in the past 30 days. <sup>24</sup> Among those

59.5

63.6

55.9\*

 $\textbf{NOTE:} \ \ \text{Data are weighted to be nationally representative of all students aged } 13-15\ \text{years who are enrolled in school.} \ \ \text{Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.}$ 

inside enclosed public places

outdoor public places

Students who favored prohibiting smoking at

<sup>\*</sup>Gender comparisons are significant at p < 0.05.